

Collaborative Research on the Northeast Water Polynya: NEWP92
Hydrographic Data Report
USCGC *Polar Sea* Cruise, July 15 - August 15, 1992

D.W.R. Wallace¹
T.S. Hopkins²
W.J. Behrens¹
J. Deming³
C. Kinder²
W.O. Smith⁴
Z. Top⁵
I.D. Walsh⁶

¹Oceanographic and Atmospheric Sciences Division, Department of Applied Science,
Brookhaven National Laboratory, Upton, NY 11973

²Department of Marine, Earth and Atmospheric Science, North Carolina State University,
Raleigh, NC 27695

³School of Oceanography, University of Washington, WB10, Seattle, WA 98195

⁴Graduate Program in Ecology, University of Tennessee, Knoxville, TN 37996

⁵University of Miami, Rosenstiel School of Marine and Atmospheric Sciences,
4600 Rickenbacker Causeway, Miami, FL 33149

⁶Department of Oceanography, Texas A&M University, College Station, TX 77843

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe any privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor, or subcontractor thereof.

Printed in the United States of America
Available From
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

NTIS price codes:
Printed Copy: A06; Microfiche Copy: A01

Sequence of Data Report Contents

- Cruise description; Methods; Acknowledgements; References
- Station and Cast Listing
- Station Map
- Section Plots of Properties
- Vertical Profiles of CTD Properties
- CTD and Bottle Data Listings

Introduction

The Northeast Water Polynya (NEW) off the northeast coast of Greenland was the focus of two cruises aboard the USCGC *Polar Sea* during the summers of 1992 and 1993. The cruises were supported by the National Science Foundation Arctic Systems Science (ARCSS) program and were part of the Arctic Ocean Science Board's International Arctic Polynya Program. The Polar Sea cruises were designed as multidisciplinary studies to test hypotheses about the mechanisms of heat, water and carbon flow within and beyond the boundaries of the polynya. Preliminary results of the 1992 study have been described elsewhere (NEWATER, 1993). A collection of papers arising from the 1992 cruise have been published in a Special Section of the Journal of Geophysical Research (Overland et al., 1995).

This data report presents the hydrographic and basic chemical observations made from CTD/Rosette casts during the 1992 cruise. The station positions cruise are plotted in Figure 1. Also included in the report are selected section plots and vertical profiles. A total of 130 CTD casts were made during the cruise, measuring pressure, temperature, conductivity, dissolved oxygen, fluorescence and light transmission. Discrete samples were collected in 10-liter, rosette-mounted, Niskin bottles and analyzed, from most casts, for: salinity, dissolved nutrients, dissolved oxygen, anthropogenic halocarbons (e.g., Freon gases), pigments, particulate organic carbon and nitrogen. Suspended particulate matter was analyzed at selected stations and these data were used to calibrate the CTD-transmissometer. Samples were collected from selected stations and depths for tritium and helium analyses, carbonate chemistry, as well as for measurements of bacterial abundance.

All the data listed in this report are currently stored in a remotely-accessible database at Brookhaven National Laboratory, access to which can be arranged on request (contact: D. Wallace). In addition, the entire NEWP92 and NEWP93 dataset is to be published in CD-ROM format by the National Snow and Ice Data Center, University of Colorado, Boulder, CO. The hydrographic data will also be submitted to NOAA's National Oceanographic Data Center.

Description of Methods:

Header Data:

Ice cover refers to bridge observations made by Coast Guard personnel at or close to the time of sampling. Specifically, ice cover at any particular station has been estimated from linear interpolation of the hourly bridge observations which were initially recorded in tenths. It should be noted that the ice cover in this region was often highly variable in time, as well as in space.

CTD Data:

CTD/Rosette Sampling. CTD data were collected using a Neil Brown Instrument Systems Mark III CTD underwater unit with a model 1150 deck data terminal. The CTD was equipped with a Beckman oxygen sensor and interfaced with a SeaTech fluorometer, and a 25cm path-length SeaTech transmissometer.

All underwater instrumentation was mounted on a General Oceanics rosette multi-bottle sampler, model 1015-12. The rosette was equipped with 11 x 10-liter Niskin bottles, and a single 5-liter bottom-tripped bottle. The bottom-trip switch was triggered when a weight suspended from a lanyard one meter beneath the rosette contacted the bottom.

CTD data were collected and displayed in real-time on a PC-486 computer using Oceansoft I software from EG&G Marine Instruments.

CTD Data Processing. The raw CTD data were transferred to a MicroVAX for processing. Large spikes were removed using a gradient check and 0.1 dbar averages were calculated. For each cast, data were extrapolated to the bottom depth. Any obvious large-amplitude spikes that remained in the raw data variables (pressure, temperature, conductivity, oxygen current, oxygen temperature, fluorometer and transmissometer) were edited and replaced by interpolation. Additionally, any missing values were filled in by linear interpolation.

To reduce the spikes caused in the calculation of salinity, a three-point moving average was first performed on conductivity to create a smooth signal. A lag of .4 was then applied to conductivity to better match the conductivity sensor with the temperature sensor. After final calibrations were applied to the CTD data, derived variables (depth, potential temp, salinity, oxygen in %-saturation and $\mu\text{mole kg}^{-1}$, c_p , SPM, sigma-t, sigma-p, sigma-1000, sigma-theta, Brunt-Vaisalla frequency, integrated density, steric height and freezing point) were calculated and then averaged using a pressure-centering procedure.

CTD Data Calibration. CTD data were calibrated solely with in situ methods and did not employ the use of pre- or post-cruise laboratory calibrations.

1. Pressure. A pressure offset of -0.5 dbar was applied to each CTD cast in its entirety. This correction accounted for the difference between the true and actual distance measured by the CTD pressure sensor at the surface of the water.
2. Temperature. CTD temperature was calibrated based on a linear regression with two in situ SIS reversing thermometers. A reiterative procedure produced the following relationship:

$$\text{Corr T} = .996306 * \text{CTD T} + .002380,$$

which had a standard deviation of 0.007°C.

3. Conductivity. Final conductivity corrections were derived from a reiterative linear regression of upcast CTD conductivity and in situ bottle conductivity (inverted from salinity using corrected pressure and temperature). A pressure edit greater than 30 was used to eliminate bottles taken in the chlorophyll max layer. Time-series plots of water sample data minus CTD upcast conductivity differences were constructed to determine station subgroups. Three groups were identified, giving the following slope and bias terms, and their respective standard deviations:

$$\text{CTD 1 - 89} \quad \text{Corr C} = .998553 * \text{CTD C} + .063721 \quad \text{Std. dev. } 0.007$$

$$\text{CTD 90 - 97} \quad \text{Corr C} = .953416 * \text{CTD C} + .032105 \quad \text{Std. dev. } 0.011$$

$$\text{CTD 98 - 130} \quad \text{Corr C} = .933751 * \text{CTD C} + .033966 \quad \text{Std. dev. } 0.005$$

4. Oxygen. Using the corrected pressure, salinity and temperature values, CTD values of oxygen were calculated from the upcast oxygen current and oxygen temperature sensors, following the algorithm of Owens and Millard (1985). Final oxygen coefficients were derived from reiterative linear regression of CTD upcast oxygen and in situ water sample oxygen, both in units of percent saturation. The same pressure edit criterion ($P > 30$) was used as stated above. Time-series plots yielded two groups, giving the following terms:

$$\text{CTD 1-100 Corr Ox \%} = .991460 * \text{CTD Ox \%} + .068328 \quad \text{Std. dev. 0.0099}$$

$$\text{CTD 101-130 Corr Ox \%} = .851967 * \text{CTD Ox \%} + .139818 \quad \text{Std. dev. 0.0097}$$

5. Transmissometer. Transmissometer data were calibrated by correlation with gravimetric analysis of suspended particulate loads. Between 1 and 10 liters of water were vacuum-filtered in-line from 10l water bottles through pre-weighed 0.4 μm Poretics PCTE membrane filters. The filters were rinsed in a laminar flow hood ten times with distilled, deionized and 0.4 μm filtered water, then air dried in the hood. At least one filter was put through this process during each cast without exposing it to seawater to establish an average blank value for methodological corrections. On deck transmissometer voltages were monitored throughout the cruise to assure clear optical pathways. Air calibrations of the transmissometer used were made in the lab at the beginning and end of the cruise to correct for any decay of the LED beam intensity. Readings with the light path blocked and unblocked were made through the CTD electronics prior to several casts to test for any voltage losses through the CTD. Percent transmission was recorded through the CTD and converted to a total beam attenuation coefficient (c) using the equation:

$$V/5 = T = e^{-cz}, \quad (1)$$

where V is the instrument voltage output, 5 is the maximum voltage of the transmissometer output, T is percent transmission, c is the beam attenuation coefficients with units of m^{-1} , and z is the optical pathlength in meters. Thermal hysteresis was not corrected for because the thermal variability was small ($<5^\circ\text{C}$) between the on-deck temperature and through the water column, and little or no hysteresis was apparent from comparison of up and down traces.

The beam attenuation coefficient in natural seawater is a summation of the beam attenuation coefficients for seawater (c_w , "yellow matter" (c_y), and particles (c_p) (Pak et al. 1988):

$$C = C_w + C_y + C_p. \quad (2)$$

The contribution of C_w is constant and is set at the factory to equal 0.364 m^{-1} in particle-free water for this instrument. At 660 nm, C_y is assumed constant and negligible (Bricaud et al., 1981; Gardner et al., 1993). To obtain C_p , c was calculated from the transmission values. The minimum value, c_{min} , for the cruise was subtracted from all c values to yield a pre-adjusted C_p (beam attenuation due to particles alone). Linear fits were performed between the pre-adjusted C_p values and the suspended particulate matter (SPM) concentrations obtained from the water filtered from the rosette water bottles. The data were split at a $C_p = 0.25$, essentially dividing the near surface data from the rest of the water-column. The x-intercept value from the $C_p < 25$ relationship was subtracted from the pre-adjusted C_p to obtain a C_p that was zero for a particle mass

concentration of zero. The two linear fits were then equilibrated at $c_p = 0.453$ to yield a single relationship between c_p and SPM concentration for the entire range of transmissometer data.

Bottle Data

BNL_ID: Each Niskin bottle closed during the cruise was assigned a unique identifier number that was incremented throughout the cruise; this same number was used to identify all sub-samples collected from that particular Niskin. This 'BNL_ID' number was subsequently used to merge data arising from various analytical procedures into a single database.

Dissolved Oxygen: Sub-samples for Winkler titrations were drawn into ~125 ml flasks immediately after the rosette was brought on deck and samples had been withdrawn for chlorofluorocarbon analyses. These samples were analyzed following the methodology described by Carpenter (1965). Subtle changes to methodology and calibration as described by Culberson (1991) were implemented, so that the oxygen data meet the precision and accuracy guidelines of the World Ocean Circulation Experiment (WOCE, 1991): namely accuracy <1% and precision ~0.1%.

Dissolved Nutrients: Sub-samples of ~60 ml were drawn from the Niskin bottles within 10-20 minutes of the cast being complete. Phosphate, silicate, nitrate, nitrite and ammonium were measured as soon as possible after sample collection (usually within a few hours) using a Technicon Autoanalyzer II, following standard colorimetric methods. The methods used have been described in Whitley et al. (1981) with the exception of the phosphate determination which used the hydrazine reductant method described in Gordon et al. (1992). Standards were analyzed with each batch of samples in order to compensate for instrument response drift. Standards were prepared in both distilled, deionized water and low-nutrient surface seawater to determine the salt-effect on colorimeter response. The wash water was distilled, deionized water.

Total Dissolved Inorganic Carbon and Total Alkalinity: Samples were collected in 500 ml ground-glass stoppered bottles, to which 200 μ l of 50%-saturated $HgCl_2$, was added. A headspace was introduced to allow for expansion, and the bottle was then tightly stoppered. The stopper was coated with Apiezon grease, and held firmly in place with elastic-bands. Samples were kept cool (~5°C) following collection and prior to analysis. Subsequently they were analyzed in the laboratory at Brookhaven National Laboratory for total dissolved inorganic carbon by coulometric titration (using a SOMMA system: Johnson and Wallace, 1992; Johnson et al., 1993). Small corrections were applied to account for the loss of CO_2 gas to the headspace, and for the dilution of the sample due to addition of $HgCl_2$. Total alkalinity analyses were performed subsequently on the same samples at the Woods Hole Oceanographic Institution (Dr. Catherine Goyet's laboratory) using potentiometric titration in a closed, glass cell.

Pigments: Subsamples (~280 ml) were filtered through Gelman GF/F glass fiber filters (~0.7 μ m nominal pore size). The filters were sonicated (on ice, in darkness) for 10 minutes in glass 15 ml centrifuge tubes together with 10 ml of 90% acetone. The samples were extracted in the dark for an additional 15 minutes and read using a Turner Designs Model 10 fluorometer before and after acidification (Smith and Nelson, 1990). The fluorometer was calibrated at the beginning and end

of the cruise using a commercially prepared chlorophyll standard.

Particulate Organic Carbon and Nitrogen: POC and PON were quantified by filtering samples onto precombusted (450°C for 2 hours) GF/F Whatman glass fiber filters, placing them in combusted glass vials, drying the filters at 60° C, and analyzing them on a Carlo-Erba model EA1108 elemental analyzer after high-temperature pyrolysis.

Tritium and Helium: For tritium analysis, 1-liter water samples were drawn from the Niskins into glass bottles, which had been previously filled with argon gas. These bottles were capped with a minimum of head-space and transferred to the laboratory. In the laboratory the samples were transferred to 1-liter Corning 1724 glass flasks in a vacuum line, degassed, sealed, and placed in a storage freezer at -20°C. After about 8 months, accumulated ³He in the flask was measured in a MAP215-50 mass-spectrometer and the tritium concentration was calculated using calibrated air aliquots as standards, and appropriate correction factors. In addition, a set of NBS standards prepared to have 1 - 2 TU were measured as a check on accuracy. The reproducibility of the NBS standards and duplicate samples was 0.007 TU. Analytical precision was ~2.5% of the measurement or ~0.007 TU, whichever was greater (1 TU = 1 ³He/10¹⁸H). Measurements were decay-corrected to the time of sampling.

For helium analysis, 50 g water samples were sealed in refrigeration-grade copper tubes by clamping, and transferred to the laboratory. In the laboratory, water samples were processed in a vacuum line to seal the helium-neon fraction of the dissolved gas into Corning 1724 glass ampoules. These were then re-processed in the mass-spectrometer inlet line to separate helium from neon, and the helium fraction was admitted into the MAP 215-50 mass-spectrometer. ³He and ⁴He beams were measured separately, and calibrated with standard air aliquots. The ³He ratio anomaly was expressed as:

$$\delta^3\text{He} = [[(^3\text{He}/^4\text{He})_{\text{sx}} / (^3\text{He}/^4\text{He})_{\text{std}}] - 1] \times 100,$$

where sx and std refer to the sample and the atmospheric standard, respectively. ⁴He measurements were converted to concentrations by peak height comparison, and were expressed in units of nanomoles per kg (seawater). The precision in $\delta^3\text{He}$ is $\pm 0.2\%$; in ⁴He concentration, it was $\pm 1.0\%$.

Bacterial Abundance. Water samples were collected aseptically directly from the Niskin bottles immediately after arrival of the CTD on deck. Volumes of 10 ml were fixed in 0.2- μm filtered 2% formaldehyde and stored at 4°C in the dark until processed for counting. We used a dual staining procedure, reported by Deming et al. (1995), to enumerate bacteria. Samples were stained with acridine orange (AO), according to Hobbie et al. (1977), and gently filtered onto a 0.2- μm black Nucleopore filter, followed by DAPI stain (0.001%) according to Porter and Feig (1980). The slide preparations were viewed using a Zeiss epifluorescence microscope. Individual bacteria in 20 randomly selected fields (a minimum of 200 bacteria per slide) were counted routinely using standard optical filters for AO; switching to optical filters for DAPI allowed confirmation that AO-fluorescing particles were microorganisms.

Units: All concentration data are reported in units of per kg (seawater), with the exception of bacterial abundance which is in units of cells per ml.

Acknowledgments

This work was supported by grants from the National Science Foundation, Division of Polar Programs.

Nutrient analyses were performed by Dennis Guffey (Texas A&M University) and Robert Ramirez (BNL). Oxygen analyses and other chemical sampling were performed by Craig Neill (BNL). Anne Close (Bermuda Biological Laboratory) and Shelly Carpenter (Univ. Washington) assisted with the chemical sample collection. Rick Wilke (BNL) assisted with the final nutrient data processing. Tish Yager (Univ. Washington) performed all of the inorganic carbon and alkalinity measurements. Ken Johnson provided guidance and assistance with the TCO₂ measurements. Catherine Goyet (Woods Hole Oceanographic Institution) very kindly allowed us to make the alkalinity measurements in her laboratory. Jan Gaylord and Shelly Carpenter (Univ. Washington) performed the enumeration of bacteria. Carin Ashjian and Peter Minnett (BNL) extracted the appropriate bridge ice-observations for the CTD casts. CTD operations were assisted by Dave Voegel (TAMU) and John Smithhauser (U. Alaska). Figures and final report preparation assistance from Yan Shi (BNL) is appreciated.

References

Bricaud, A., A. Morel and L. Prieur (1981) Absorption by dissolved organic matter of the sea (yellow substance) in the UV and visible domains. *Limnol. Oceanogr.* 26: 43-53.

Carpenter, J.H. (1965) The Chesapeake Bay Institute technique for the Winkler dissolved oxygen method. *Limnol. Oceanogr.* 10: 141-143.

Culberson, C.H. (1991) WHP Operations and Methods: Dissolved Oxygen. In WOCE Operations Manual, Volume 3: The Observational Programme. Part 3.1.3. WHP Operations and Methods. WHP Office Report WHPO 91-1; WOCE Report No. 68/91. Woods Hole, MA. July 1991.

Doming, J.W., A.-L. Reysenbach, S.A. Macko, and C.R. Smith (1995) Evidence for the microbial basis of a chemoautotrophic invertebrate community at a whale fall on the deep seafloor: bone-colonizing bacteria and invertebrate endosymbionts. *J. Microscopy Res. Tech.* (in press).

Gardner, W.D., I.D. Walsh and M.J. Richardson (1993) Biophysical forcing of particle production and distribution during a spring bloom in the North Atlantic. *Deep-Sea Res.* 40: 171-195.

Gordon, L.I., J.C. Jennings, A.A. Ross and J.M. Krest (1992) A suggested protocol for continuous flow automated analysis of seawater nutrients. WOCE Hydrographic Program Office, Methods Manual (In Prep.), Woods Hole, MA.

Hobbie, J.E., R.J. Daley, and S. Jasper (1977) Use of Nuclepore filters for counting bacteria by fluorescence microscopy. *Appl. Environ. Microbiol.* 33: 1225-1228.

Johnson, K.M. and D.W.R. Wallace (1992) The Single-Operator Multiparameter Metabolic Analyzer for Total Carbon Dioxide with Coulometric Detection. DOE Research Summary No. 19, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Johnson, K.M., K.D. Wills, D.B. Butler, W.K. Johnson and C.S. Wong (1993) Coulometric total carbon dioxide analysis for marine studies: maximizing the performance of an automated gas extraction system and Coulometric detector. *Marine Chemistry*, 44, 167-187.

NEWATER Science Steering Committee and Principal Investigators (1993) Northeast Water Polynya: Polar Sea Cruise Results. *EOS*, 74(16): 185, 195-196.

Overland, J.E., T.B. Curtin and W.O. Smith, Jr. (1995) Preface to Special Section: Leads and Polynyas. *J. Geophys. Res.*, 100, 4267-4268.

Pak, H., D.A. Kiefer and J.C. Kitchen (1988) Meridional variations in the concentration of chlorophyll and microparticles in the North Pacific Ocean. *Deep-Sea Res.* 35: 1151-1171.

Porter, K.G. and Y.S. Feig (1980) The use of DAPI for identifying and counting aquatic microflora. *Limnol. Oceanogr.* 25, 943-948.

Smith, W.O., Jr. and D.M. Nelson (1990) Primary productivity and new production in the Weddell Sea marginal ice zone during austral spring and autumn, *Limnol. Oceanogr.*, 35, 809-821.

Whitledge, T., S. Malloy, C. Patton and C.D. Wirrick (1981) Automated nutrient analysis in seawater. Brookhaven National Laboratory Technical Report, BNL 51398.

WOCE (1991) WOCE Operations Manual, Volume 3: The Observational Programme. Part 3.1.2. Requirements for WHP Data Reporting. WHP Office Report WHPO 90-1; WOCE Report No. 67/91. Woods Hole, MA. July 1991.

NEWP92 STATION LIST

| STATION | CAST | CTD-NO | DATE | TIME | LATITUDE | LONGITUDE | SONIC DEPTH METERS |
|---------|------|--------|---------|-------|----------|-----------|--------------------|
| | | | GMT | GMT | DEG. | DEG. | |
| 1 | 1 | 1 | 7/18/92 | 08:42 | 76.526 | -6.907 | 1343 |
| 1 | 2 | 2 | 7/18/92 | 11:15 | 76.512 | -6.966 | 1304 |
| 2 | 3 | 3 | 7/19/92 | 00:28 | 77.155 | -10.481 | 488 |
| 2 | 4 | 4 | 7/19/92 | 01:49 | 77.160 | -10.447 | 476 |
| 2 | 5 | 5 | 7/19/92 | 03:01 | 77.160 | -10.406 | 472 |
| 2 | 6 | 6 | 7/19/92 | 11:21 | 77.150 | -10.449 | 487 |
| 3 | 7 | 7 | 7/20/92 | 12:43 | 78.606 | -13.487 | 142 |
| 3 | 8 | 8 | 7/20/92 | 14:17 | 78.609 | -13.508 | 139 |
| 3 | 9 | 9 | 7/20/92 | 15:28 | 78.616 | -13.528 | 137 |
| 3 | 10 | 10 | 7/20/92 | 19:49 | 78.645 | -13.494 | 168 |
| 4 | 11 | 11 | 7/21/92 | 13:55 | 80.398 | -14.243 | 337 |
| 4 | 12 | 12 | 7/21/92 | 15:11 | 80.410 | -14.208 | 333 |
| 4 | 13 | 13 | 7/21/92 | 16:29 | 80.421 | -14.156 | 330 |
| 5 | 14 | 14 | 7/22/92 | 13:12 | 80.652 | -11.269 | 238 |
| 6 | 15 | 15 | 7/22/92 | 15:51 | 80.649 | -12.169 | 260 |
| 7 | 16 | 16 | 7/22/92 | 19:57 | 80.648 | -13.082 | 222 |
| 8 | 17 | 17 | 7/24/92 | 02:29 | 80.532 | -14.661 | 208 |
| 9 | 18 | 18 | 7/24/92 | 05:22 | 80.483 | -14.488 | 324 |
| 10 | 19 | 19 | 7/24/92 | 07:03 | 80.440 | -14.422 | 301 |
| 11 | 20 | 20 | 7/24/92 | 08:16 | 80.369 | -14.088 | 314 |
| 12 | 21 | 21 | 7/24/92 | 10:33 | 80.307 | -13.797 | 286 |
| 13 | 22 | 22 | 7/24/92 | 12:22 | 80.238 | -13.617 | 175 |
| 14 | 23 | 23 | 7/24/92 | 13:39 | 80.188 | -13.454 | 185 |
| 15 | 24 | 24 | 7/24/92 | 16:01 | 80.119 | -13.159 | 75 |
| 16 | 25 | 25 | 7/25/92 | 08:33 | 80.275 | -11.029 | 199 |
| 17 | 26 | 26 | 7/25/92 | 11:13 | 80.189 | -10.923 | 121 |
| 18 | 27 | 27 | 7/25/92 | 13:44 | 80.338 | -11.026 | 253 |
| 19 | 28 | 28 | 7/25/92 | 15:31 | 80.409 | -11.031 | 315 |
| 20 | 29 | 29 | 7/25/92 | 17:43 | 80.477 | -10.897 | 290 |
| 21 | 30 | 30 | 7/25/92 | 19:44 | 80.539 | -10.981 | 257 |
| 22 | 31 | 31 | 7/25/92 | 21:16 | 80.609 | -10.975 | 267 |
| 23 | 32 | 32 | 7/25/92 | 22:17 | 80.671 | -10.956 | 181 |
| 24 | 33 | 33 | 7/25/92 | 23:41 | 80.722 | -10.938 | 111 |
| 25 | 34 | 34 | 7/26/92 | 08:37 | 80.443 | -13.332 | 288 |
| 25 | 35 | 35 | 7/26/92 | 14:12 | 80.450 | -13.349 | 306 |
| 25 | 36 | 36 | 7/26/92 | 20:58 | 80.455 | -13.401 | 305 |
| 25 | 37 | 37 | 7/27/92 | 03:16 | 80.454 | -13.328 | 301 |
| 25 | 38 | 38 | 7/27/92 | 03:51 | 80.458 | -13.302 | 301 |
| 25 | 39 | 39 | 7/27/92 | 04:43 | 80.464 | -13.272 | 302 |

| | | | | | | | |
|----|----|----|---------|-------|--------|---------|-----|
| 26 | 44 | 44 | 7/28/92 | 12:15 | 80.337 | -14.466 | 344 |
| 27 | 45 | 45 | 7/28/92 | 13:50 | 80.196 | -15.283 | 271 |
| 28 | 46 | 46 | 7/28/92 | 16:49 | 80.014 | -15.944 | 418 |
| 29 | 47 | 47 | 7/28/92 | 22:16 | 80.093 | -16.105 | 187 |
| 30 | 48 | 48 | 7/29/92 | 00:20 | 80.077 | -15.811 | 350 |
| 30 | 49 | 49 | 7/29/92 | 03:36 | 80.075 | -15.813 | 354 |
| 31 | 50 | 50 | 7/29/92 | 04:56 | 80.049 | -15.505 | 275 |
| 32 | 51 | 51 | 7/29/92 | 10:08 | 80.025 | -15.137 | 234 |
| 33 | 52 | 52 | 7/29/92 | 11:32 | 79.985 | -14.789 | 149 |
| 34 | 53 | 53 | 7/29/92 | 13:34 | 79.939 | -14.213 | 85 |
| 35 | 54 | 54 | 7/29/92 | 17:10 | 80.020 | -13.426 | 83 |
| 36 | 55 | 55 | 7/29/92 | 21:26 | 80.092 | -13.940 | 154 |
| 37 | 56 | 56 | 7/29/92 | 23:34 | 80.168 | -14.368 | 248 |
| 38 | 57 | 57 | 7/30/92 | 00:51 | 80.214 | -14.612 | 332 |
| 39 | 58 | 58 | 7/30/92 | 02:12 | 80.262 | -14.911 | 380 |
| 40 | 59 | 59 | 7/30/92 | 04:56 | 80.321 | -15.228 | 272 |
| 41 | 60 | 60 | 7/30/92 | 06:41 | 80.383 | -15.583 | 93 |
| 42 | 61 | 61 | 7/30/92 | 10:09 | 80.020 | -15.988 | 437 |
| 42 | 62 | 62 | 7/30/92 | 11:38 | 80.031 | -16.007 | 388 |
| 42 | 63 | 63 | 7/30/92 | 16:39 | 80.019 | -15.971 | 450 |
| 42 | 64 | 64 | 7/30/92 | 18:21 | 80.029 | -15.986 | 450 |
| 43 | 65 | 65 | 7/31/92 | 06:44 | 80.314 | -9.670 | 313 |
| 43 | 66 | 66 | 7/31/92 | 09:11 | 80.312 | -9.698 | 312 |
| 43 | 67 | 67 | 7/31/92 | 15:42 | 80.311 | -9.549 | 308 |
| 43 | 68 | 68 | 7/31/92 | 21:28 | 80.306 | -9.528 | 309 |
| 43 | 69 | 69 | 8/01/92 | 01:37 | 80.312 | -9.191 | 305 |
| 43 | 70 | 70 | 8/01/92 | 06:16 | 80.312 | -9.657 | 312 |
| 43 | 71 | 71 | 8/01/92 | 09:23 | 80.265 | -9.656 | 303 |
| 43 | 72 | 72 | 8/01/92 | 14:12 | 80.339 | -9.502 | 306 |
| 43 | 73 | 73 | 8/01/92 | 15:26 | 80.319 | -9.686 | 315 |
| 43 | 74 | 74 | 8/01/92 | 20:53 | 80.297 | -9.426 | 314 |
| 43 | 75 | 75 | 8/01/92 | 21:57 | 80.284 | -9.441 | 310 |
| 43 | 76 | 76 | 8/02/92 | 03:27 | 80.321 | -9.696 | 316 |
| 44 | 77 | 77 | 8/02/92 | 16:21 | 80.744 | -7.643 | 175 |
| 45 | 78 | 78 | 8/02/92 | 18:46 | 80.731 | -7.326 | 276 |
| 46 | 79 | 79 | 8/02/92 | 22:12 | 80.733 | -7.154 | 356 |
| 47 | 80 | 80 | 8/03/92 | 10:29 | 80.675 | -8.245 | 69 |
| 48 | 81 | 81 | 8/04/92 | 13:42 | 80.720 | -8.819 | 58 |
| 48 | 82 | 82 | 8/04/92 | 15:34 | 80.783 | -8.825 | 61 |
| 49 | 83 | 83 | 8/05/92 | 10:08 | 80.770 | -8.181 | 76 |
| 49 | 84 | 84 | 8/05/92 | 11:19 | 80.742 | -8.238 | 75 |
| 50 | 85 | 85 | 8/06/92 | 07:56 | 80.606 | -7.911 | 101 |
| 51 | 86 | 86 | 8/06/92 | 10:59 | 80.487 | -8.219 | 249 |
| 52 | 87 | 87 | 8/06/92 | 12:59 | 80.378 | -8.394 | 255 |
| 53 | 88 | 88 | 8/06/92 | 15:14 | 80.282 | -8.696 | 292 |

| | | | | | | | |
|----|-----|-----|---------|-------|--------|---------|------|
| 57 | 93 | 93 | 8/07/92 | 08:49 | 80.309 | -8.555 | 283 |
| 58 | 94 | 94 | 8/07/92 | 23:08 | 80.444 | -13.316 | 285 |
| 58 | 95 | 95 | 8/08/92 | 00:45 | 80.448 | -13.266 | 292 |
| 59 | 96 | 96 | 8/08/92 | 08:05 | 80.012 | -15.930 | 425 |
| 60 | 97 | 97 | 8/08/92 | 10:33 | 79.833 | -16.164 | 236 |
| 61 | 98 | 98 | 8/08/92 | 12:07 | 79.757 | -15.574 | 211 |
| 62 | 99 | 99 | 8/08/92 | 15:58 | 79.628 | -15.977 | 200 |
| 62 | 100 | 100 | 8/08/92 | 17:15 | 79.629 | -15.969 | 196 |
| 63 | 101 | 101 | 8/09/92 | 02:41 | 79.157 | -13.769 | 126 |
| 63 | 102 | 102 | 8/09/92 | 04:18 | 79.154 | -13.807 | 123 |
| 64 | 103 | 103 | 8/09/92 | 12:38 | 79.022 | -12.050 | 216 |
| 64 | 104 | 104 | 8/09/92 | 14:05 | 79.020 | -12.026 | 219 |
| 64 | 105 | 105 | 8/09/92 | 18:16 | 79.008 | -11.975 | 220 |
| 65 | 106 | 106 | 8/10/92 | 08:46 | 78.314 | -16.743 | 525 |
| 65 | 107 | 107 | 8/10/92 | 11:24 | 78.312 | -16.768 | 512 |
| 65 | 108 | 108 | 8/10/92 | 13:04 | 78.315 | -16.783 | 515 |
| 66 | 109 | 109 | 8/11/92 | 06:44 | 77.635 | -11.374 | 271 |
| 67 | 110 | 110 | 8/11/92 | 08:45 | 77.572 | -11.688 | 440 |
| 68 | 111 | 111 | 8/11/92 | 10:56 | 77.460 | -12.044 | 490 |
| 69 | 112 | 112 | 8/11/92 | 13:52 | 77.351 | -12.326 | 321 |
| 70 | 113 | 113 | 8/11/92 | 17:05 | 77.235 | -12.669 | 261 |
| 71 | 114 | 114 | 8/11/92 | 21:49 | 77.341 | -11.854 | 414 |
| 72 | 115 | 115 | 8/12/92 | 01:04 | 77.149 | -10.527 | 486 |
| 72 | 116 | 116 | 8/12/92 | 05:12 | 77.137 | -10.547 | 490 |
| 72 | 117 | 117 | 8/12/92 | 07:04 | 77.153 | -10.504 | 490 |
| 72 | 120 | 120 | 8/13/92 | 09:17 | 77.163 | -10.535 | 478 |
| 72 | 121 | 121 | 8/13/92 | 11:08 | 77.161 | -10.551 | 485 |
| 73 | 118 | 118 | 8/12/92 | 18:39 | 77.282 | -10.051 | 350 |
| 74 | 119 | 119 | 8/13/92 | 05:41 | 77.001 | -10.910 | 385 |
| 75 | 122 | 122 | 8/13/92 | 18:30 | 76.888 | -8.848 | 363 |
| 75 | 123 | 123 | 8/13/92 | 20:24 | 76.888 | -8.851 | 363 |
| 76 | 124 | 124 | 8/14/92 | 03:22 | 77.177 | -7.995 | 287 |
| 77 | 125 | 125 | 8/14/92 | 06:45 | 77.109 | -7.170 | 266 |
| 78 | 126 | 126 | 8/14/92 | 09:06 | 76.988 | -6.576 | 273 |
| 79 | 127 | 127 | 8/14/92 | 10:45 | 76.948 | -6.291 | 697 |
| 80 | 128 | 128 | 8/14/92 | 12:41 | 76.919 | -6.065 | 1007 |
| 81 | 129 | 129 | 8/14/92 | 15:04 | 76.850 | -5.562 | 1480 |
| 81 | 130 | 130 | 8/14/92 | 16:50 | 76.834 | -5.561 | 1490 |